1 stepped pressure equilibrium code : fe00aa

Contents

1	stepped pressure equilibrium code : fe00aa	1
	1.1 outline	1

1.1 outline

- 1. Calculates finite element functions, and their derivatives, on radial sub-sub-grid. The quantities calculated are the $\varphi_{l,p}$ for l=0,1 and p=0, Nofe which are stored in array felr(0:1,0:Nofe,0:1,1:Nj) (which is allocated in readin).
- 2. This routine is only called once, from xspech.
- 3. Should check that $\varphi_{1,p}(x) = (-1)^p \varphi_{0,p}(1-x)$ is satisfied.
- 4. Radial quadrature is performed using Gaussian quadrature. The resolution of the approximation is controlled by Iquad, and the weights and abscissae are returned by NAG routine D01BCF. At present, only itype=0 (Gauss-Legrendre) and a=0, b=1, c=0 and d=0 are supplied as input to D01BCF.
- 5. The array felr(0:1,0:Nofe,0:1,1:Iquad) is constructed as follows:
 - (a) the first argument is the left-right index: $felr(0,p,0,j) = \varphi_{0,p}(x_j)$ and $felr(1,p,0,j) = \varphi_{1,p}(x_j)$;
 - (b) the second argument is the order of the basis function:
 - (c) the third argument is the derivative: $felr(0,p,1,j) = \varphi'_{L,p}(x_j);$
 - (d) and the fourth argument indicates the radial sub-sub-grid index.
- 6. Note that the derivatives with respect to x are returned, where $x = (s s_{l,i-1})/\Delta s_l$ and $\Delta s_l = (s_{l,i} s_{l,i-1})$. The radial scaling factor Δs_l needs to be included elsewhere.
- 7. The radial basis functions used are:
 - Nofe=0: linear basis functions;
 - Nofe=1: cubic basis functions;

$$\varphi_{0,0} = 2 \quad x^3 - 3 \quad x^2 + 1
\varphi_{0,1} = \quad x^3 - 2 \quad x^2 + x
\varphi_{1,0} = -2 \quad x^3 + 3 \quad x^2
\varphi_{1,1} = \quad x^3 - x^2$$
(1)

• Nofe=2: quintic basis function;

$$\varphi_{L,0} = -6 \quad x^5 + 15 \quad x^4 - 10 \quad x^3 + 1
\varphi_{L,1} = -3 \quad x^5 + 8 \quad x^4 - 6 \quad x^3 + 1
\varphi_{L,2} = -1/2 \quad x^5 + 3/2 \quad x^4 - 3/2 \quad x^3 + 1/2 \quad x^2
\varphi_{R,0} = 6 \quad x^5 - 15 \quad x^4 + 10 \quad x^3
\varphi_{R,0} = -3 \quad x^5 + 7 \quad x^4 - 4 \quad x^3
\varphi_{R,0} = 1/2 \quad x^5 - x^4 + 1/2 \quad x^3$$
(2)

• Nofe=3: septemtic basis function;

$$\begin{array}{l} \varphi_{0,0} &=& 20 \quad x^7 \quad - \quad 70 \quad x^6 \quad + \quad 84 \quad x^5 \quad - \quad 35 \quad x^4 \\ \varphi_{0,1} &=& 10 \quad x^7 \quad - \quad 36 \quad x^6 \quad + \quad 45 \quad x^5 \quad - \quad 20 \quad x^4 \\ \varphi_{0,2} &=& 2 \quad x^7 \quad - \quad 15/2 \quad x^6 \quad + \quad 10 \quad x^5 \quad - \quad 5 \quad x^4 \\ \varphi_{0,3} &=& 1/6 \quad x^7 \quad - \quad 2/3 \quad x^6 \quad + \quad 1 \quad x^5 \quad - \quad 2/3 \quad x^4 \quad + \quad 1/6 \quad x^3 \\ \varphi_{1,0} &=& -20 \quad x^7 \quad + \quad 70 \quad x^6 \quad - \quad 84 \quad x^5 \quad + \quad 35 \quad x^4 \\ \varphi_{1,1} &=& 10 \quad x^7 \quad - \quad 34 \quad x^6 \quad + \quad 39 \quad x^5 \quad - \quad 15 \quad x^4 \\ \varphi_{1,2} &=& -2 \quad x^7 \quad + \quad 13/2 \quad x^6 \quad - \quad 7 \quad x^5 \quad + \quad 5/2 \quad x^4 \\ \varphi_{1,3} &=& 1/6 \quad x^7 \quad - \quad 1/2 \quad x^6 \quad + \quad 1/2 \quad x^5 \quad - \quad 1/6 \quad x^4 \end{array}$$

(3)

• otherwise there is an error;

fe00aa.h last modified on 2012-09-21;